

A Roadmap For Meson Spectroscopy

M.G. Olsson

Physics Department, University of Wisconsin, Madison, WI 53706

An efficient classification of light quark meson states is discussed based on the dominance of angular and radial quark excitation. A synthesis of Regge and quark dynamics allows a natural unification of light quark-antiquark spectroscopies and indicates the states that fall outside this category such as molecules, hybrids, and glueballs.

I. INTRODUCTION

The rapidly evolving field of hadron spectroscopy is increasingly motivated by firm expectations of seeing new types of matter[1]. The “ordinary” mesons are thought to be confined quark-antiquark pairs in spin singlet/triplet states and with orbital and radial excitation. Non-perturbative QCD in addition predicts that for each ordinary meson there are higher lying states with excited gluonic energy (hybrids) and that there should also be mesons composed essentially of pure glue energy (glueballs). Further complicating the experimental situation are possible four quark mesons (molecules) or dynamically generated resonances not having any clear relation to the quark picture.

In the most recent Particle Data Group report [2] there are about 100 well established meson states composed of light quarks (u, d, s) and glue. Of these, 56 have been assigned into constituent quark multiplets [2]. Since the vast majority of the observed meson states lie in energy below 2.5 GeV a straightforward method of separating the ordinary mesons from hybrids, glueballs etc. would be very useful. Previous work [3] in understanding the meson spectrum has either centered on detailed quark models or on the search for SU(3) flavor nonets. Both approaches are sensitive to details such as mixing between nonets or with hybrids and glueballs. Although such considerations are crucial to a complete understanding, the proposed classification is based on some simple observations and is relatively model independent.

First we draw some motivation from the well-understood heavy onia mesons composed of b and c quarks. These states are classified by the orbital angular momentum and radial excitation of the heavy quarks. The spin dependence is much smaller than the quark motion energies and therefore each state of orbital and radial excitation consists for $L > 0$ of four nearly degenerate states (e.g. χ_0, χ_1, χ_2 and h). For $L = 0$ the group is just two states. The corresponding Regge trajectories look very different when plotted conventionally as total angular momentum J vs. E^2 . Here it is evident that there is a “fundamental” orbital angular momentum trajectory $L = L(E^2)$ that breaks into a family of trajectories when the spin is turned on. Nothing spectacular happens in the case of light quarks except that the spin splittings are somewhat larger. Thus our first principle will be to consider all light quark states of the same orbital and radial excitation quantum numbers to form natural groups. For $L > 0$ and three light quarks there are expected to be sixteen members of each group (and eight members for $L = 0$ groups). Clearly, a few such groups accounts

for a lot of mesons.

For light quarks (and actually also true for heavy quark bound states) the confining interaction dominates the meson dynamics. The confining interaction closest to QCD is the QCD string/flux tube. The flux tube has all the proper QCD relativistic corrections [4] as well as the correct Regge slope of $1/2\pi a$ [5]. A string tension of $a = 0.16 \text{ GeV}^2$ yields the universal Regge slope of 1 GeV^{-2} . In addition the long range excited glue spectrum is correctly predicted[6]. The light quark QCD flux tube spectroscopy can be exactly numerically calculated[4, 6] and analytically accurately estimated [7] by,

$$E^2/2\pi a = L + 2n - 1/2, \quad (1)$$

where E is the meson excitation energy and L takes the values $0, 1, 2, \dots$ and $n = 1, 2, \dots$. We should note the factor $L + 2n$ which implies degenerate tower states of either even or odd L .

The effect of an attractive short-range interaction will be to lower E for a given L . For large L the short-range interaction, $V(r) = -k/r$, shifts [8] the Regge trajectory to smaller $E^2/(2\pi a)$ by $k/2$. For $L = 0$ this shift is somewhat larger. In Fig. 1 we show by the solid lines the Regge trajectories due to confinement alone. These trajectories given by Eq. (1) consist of the leading ($n = 1$) and evenly spaced daughter trajectories all of slope $1/2\pi a$ (or unit slope in this graph). This structure is numerically accurate for a straight string and also for a dynamically curved string [9]. The dashed curves represent mesons with both confinement and with short range (taking $k = 1$) interaction.

II. ORDINARY MESONS

We first consider the light quark mesons [2] that have been assigned to constituent quark configurations. We subdivide these according to their assigned orbital and radial quantum numbers L and n [2]. Taking $2\pi a = 1 \text{ GeV}^{-2}$ we plot $E^2/2\pi a$ for each of these mesons in Fig.1. In this calculation we have subtracted 0.1 GeV for each strange quark to roughly account for the differing quark masses.

The “fundamental” Regge plot of Fig.1 shows that all of the 56 assigned quark model states indeed fall into a few well defined groups. All of the expected states of the 1S, 2S, and 1P groups are present (32 mesons). The 1D group has currently 12 of the 16 expected and the 3S 4 of 8 states. We see that the 2P, and 1F groups each has 4 out of the expected 16. One expects the remaining




FIG. 1: The fundamental Regge plot for light quark mesons. The “data” points represent the 56 quark assignments of the PDG [2] with $2\pi a = 1 \text{ GeV}^2$. One hundred MeV has been subtracted for each strange quark in these mesons. The solid lines are the string confinement spectroscopy of Eq.(1). The dashed curves are confinement plus the short distance Coulomb term $V(r) = -k/r$ with $k = 1$ [8].

mesons in these groups to be present. We also indicate several groups expected at higher energies.

III. NEW MATTER

In the conservative meson summary table [2] there are several mesons that have no logical place in Fig 1. The $f_0(.60)$ is probably a dynamical state and the $f_0(.98)$ and $a_0(.98)$ are surely some sort of four quark/molecule state associated with the $K\bar{K}$ threshold. Much has been writ-

ten on the nature of these states [2]. Between 1 and 2 GeV there are the 1P, 2S, 1D, 3S, and 2P groups. Although there are numerous missing states in these groups there is no place for the $f_0(1.50)$ which may well be a glueball. There is also mounting evidence for two exotic mesons of $J^{PC} = 1^{-+}$ (at 1.4 and 1.6 GeV) which by their quantum numbers cannot be ordinary mesons. One of these is an excellent candidate for a hybrid meson. Considerable further speculation is certainly possible but will not be pursued here.

IV. SUMMARY

Using some general organizing concepts we outline the number, quantum numbers, and approximate energies of the expected quark-antiquark (ordinary) mesons. From orbital and radial motion dominance we find that all (56) of the mesons currently [2] assigned as constituent quark model states fall into well defined groups as shown in Fig.1. For massless (ultra-relativistic) quarks and with linear confinement the allowed eigenstates generally lie on straight Regge trajectories with parallel daughter trajectories [10]. The QCD string seems to be similar to QCD at large distances giving the simple spectrum (1). As seen in Fig. 1, the observed meson groups organize neatly into these fundamental trajectories when supplemented by the expected short range interaction. These groupings represent, in a model-independent way, what one might expect in the way of ordinary mesons. This should be a useful tool in the identification of the new matter states.

Acknowledgments

This research was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896 and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

-
- [1] Key Issues in Hadron Physics, Town Hall Meeting, Jefferson Laboratory, Dec. 1-4 2000, ArXiv:hep-ph/0012238.
 - [2] Particle Data Group, K.Hagiwara *et al.*, Phys. Rev. **D66**, 010001 (2002).
 - [3] S. Godfrey and N. Isgur, Phys. Rev. **D32**, 189 (1985); S. Godfrey and N. Isgur, Phys. Rev. **D31**, 2375 (1985); F.L. Close, “Quarks and Nuclear Forces” (Springer-Verlag, 1982).
 - [4] Collin Olson, M.G. Olsson, and Ken Williams, Phys. Rev. **D45**, 4307 (1992); N. Brambilla and G.M. Prosperi, Phys. Lett. B235, 69 (1990).
 - [5] D. LaCourse and M.G. Olsson, Phys. Rev. **D39**, 2751 (1989); M.G. Olsson and Sinisa Veseli, Phys. Rev. **D51**, 3578 (1995); Yu. Dubin, A.B. Kaidalov, and Yu. A. Simionov, Phys. Lett. **B323**, 41 (1994); **B343**, 310 (1995).
 - [6] T.J. Allen, M.G. Olsson, and Sinisa Veseli, Phys. Lett. **B434**, 110 (1998).
 - [7] T.J. Allen, C. Goebel, M.G. Olsson, and Sinisa Veseli, Phys. Rev. **D64**, 094011 (2001).
 - [8] The Coulomb shift is most simply established for large L for the classical (leading) Regge trajectory. For massless quarks and circular orbits the QCD string [4] gives $L = \pi ar^2/8$ and $E = \pi ar/2 - k/r$. Eliminating the quark separation r , squaring E and dropping the small $1/L$ term gives $E^2/2\pi a = L - k/2$.
 - [9] T.J.Allen, M.G. Olsson, and Sinisa Veseli, Phys. Rev. **D59**, 094011 (1999); Phys. Rev. **D60**, 074026 (1999).
 - [10] M.G. Olsson, Phys. Rev. **D55**, 5479 (1997).